



MASSACHUSETTS

Blue Cross Blue Shield of Massachusetts is an independent  
Licensee of the Blue Cross and Blue Shield Association

## Medical Policy

# Hyperthermic Intraperitoneal Chemotherapy for Select Intra-Abdominal and Pelvic Malignancies

### Table of Contents

- [Policy: Commercial](#)
- [Policy: Medicare](#)
- [Authorization Information](#)
- [Coding Information](#)
- [Description](#)
- [Policy History](#)
- [Information Pertaining to All Policies](#)
- [References](#)

### Policy Number: 048

BCBSA Reference Number: 2.03.07 (For Plan internal use only)

NCD/LCD: N/A

### Related Policies

None

### Policy

#### Commercial Members: Managed Care (HMO and POS), PPO, and Indemnity Medicare HMO Blue<sup>SM</sup> and Medicare PPO Blue<sup>SM</sup> Members

Cytoreductive surgery plus hyperthermic intraperitoneal chemotherapy (HIPEC) at the time of surgery may be considered **MEDICALLY NECESSARY** for the treatment of:

- Pseudomyxoma peritonei; and
- Diffuse malignant peritoneal mesothelioma.

The use of HIPEC may be considered **MEDICALLY NECESSARY** in newly diagnosed epithelial ovarian or fallopian tube cancer at the time of interval cytoreductive surgery when ALL of the following criteria are met:

- The individual has stage III disease\*;
- The individual is not eligible for primary cytoreductive surgery or surgery had been performed but was incomplete and will receive neoadjuvant chemotherapy and subsequent interval debulking surgery; and
- It is expected that complete or optimal cytoreduction can be achieved at time of the interval debulking surgery.

The use of HIPEC in all other settings to treat ovarian cancer, including but not limited to stage IIIC or IV ovarian cancer, is considered **INVESTIGATIONAL**.

Ovarian cancer staging is as follows:

- Stage I: The cancer is confined to the ovary or fallopian tube.

- Stage II: The cancer involves one or both ovaries with pelvic extension.
- \*Stage III: The cancer has spread within the abdomen.
- Stage IV: The cancer is widely spread throughout the body.

Cytoreductive surgery plus HIPEC are considered **INVESTIGATIONAL** for:

- Peritoneal carcinomatosis from colorectal cancer, gastric cancer, or endometrial cancer; and
- All other indications, including goblet cell tumors of the appendix.

## Prior Authorization Information

### Inpatient

- For services described in this policy, precertification/preauthorization **IS REQUIRED** for all products if the procedure is performed **inpatient**.

### Outpatient

- For services described in this policy, see below for products where prior authorization **might be required** if the procedure is performed **outpatient**.

	Outpatient
<b>Commercial Managed Care (HMO and POS)</b>	Prior authorization is <b>not required</b> .
<b>Commercial PPO and Indemnity</b>	Prior authorization is <b>not required</b> .
<b>Medicare HMO Blue<sup>SM</sup></b>	Prior authorization is <b>not required</b> .
<b>Medicare PPO Blue<sup>SM</sup></b>	Prior authorization is <b>not required</b> .

## CPT Codes / HCPCS Codes / ICD Codes

*Inclusion or exclusion of a code does not constitute or imply member coverage or provider reimbursement. Please refer to the member's contract benefits in effect at the time of service to determine coverage or non-coverage as it applies to an individual member.*

*Providers should report all services using the most up-to-date industry-standard procedure, revenue, and diagnosis codes, including modifiers where applicable.*

*The following codes are included below for informational purposes only; this is not an all-inclusive list.*

**The above medical necessity criteria MUST be met for the following codes to be covered for Commercial Members: Managed Care (HMO and POS), PPO, Indemnity, Medicare HMO Blue and Medicare PPO Blue:**

### CPT Codes

CPT codes:	Code Description
96547	Intraoperative hyperthermic intraperitoneal chemotherapy (HIPEC) procedure, including separate incision(s) and closure, when performed; first 60 minutes (List separately in addition to code for primary procedure)
96548	Intraoperative hyperthermic intraperitoneal chemotherapy (HIPEC) procedure, including separate incision(s) and closure, when performed; each additional 30 minutes (List separately in addition to code for primary procedure)

### ICD-10 Procedure Codes

ICD-10-PCS procedure codes:	Code Description
DWY38ZZ	Hyperthermia of Abdomen
DWY68ZZ	Hyperthermia of Pelvic Region

## DESCRIPTION

### **Pseudomyxoma Peritonei**

Pseudomyxoma peritonei is a clinicopathologic disease characterized by the production of mucinous ascites and mostly originates from epithelial neoplasms of the appendix. Appendix cancer is diagnosed in fewer than 1000 Americans each year; less than half are epithelial neoplasms.<sup>1</sup> The incidence of pseudomyxoma peritonei is estimated at 2 cases per 1 million individuals.<sup>2</sup> As mucin-producing cells of the tumor proliferate, the narrow lumen of the appendix becomes obstructed and subsequently leads to appendiceal perforation. Neoplastic cells progressively colonize the peritoneal cavity and produce copious mucin, which collects in the peritoneal cavity. Pseudomyxoma peritonei ranges from benign (disseminated peritoneal adenomucinosis) to malignant (peritoneal mucinous carcinomatosis), with some intermediate pathologic grades. Clinically, this syndrome ranges from early pseudomyxoma peritonei, usually discovered during imaging or laparotomy performed for another reason, to advanced cases with a distended abdomen, bowel obstruction, and starvation.

### **Treatment**

The conventional treatment of pseudomyxoma peritonei is surgical debulking, repeated as necessary to alleviate pressure effects. However, repeated debulking surgeries become more difficult due to progressively thickened intra-abdominal adhesions, and this treatment is palliative, leaving visible or occult disease in the peritoneal cavity.<sup>3</sup>

### **Peritoneal Carcinomatosis of Colorectal Origin**

Peritoneal dissemination develops in 10% to 15% of patients with colon cancer.

### **Treatment**

Despite the use of increasingly effective regimens of chemotherapy and biologic agents to treat advanced disease, peritoneal metastases are associated with a median survival of 6 to 7 months.

### **Peritoneal Carcinomatosis of Gastric Origin**

Peritoneal carcinomatosis is detected in more than 30% of patients with advanced gastric cancer and is a poor prognostic indicator. The median survival is 3 months, and 5-year survival is less than 1%.<sup>4</sup> Sixty percent of deaths from gastric cancer are attributed to peritoneal carcinomatosis.<sup>5</sup>

### **Treatment**

Current chemotherapy regimens are nonstandard, and peritoneal seeding is considered unresectable for a cure.<sup>6</sup>

### **Peritoneal Mesothelioma**

Malignant mesothelioma is a relatively uncommon malignancy that may arise from the mesothelial cells lining the pleura, peritoneum, pericardium, and tunica vaginalis testis. In the United States, 200 to 400 new cases of diffuse malignant peritoneal mesothelioma are registered every year, accounting for 10% to 30% of all-type mesothelioma.<sup>7</sup> Diffuse malignant peritoneal mesothelioma has traditionally been considered a rapidly lethal malignancy with limited and ineffective therapeutic options. The disease is usually diagnosed at an advanced stage and is characterized by multiple variably sized nodules throughout the abdominal cavity. As the disease progresses, the nodules become confluent to form plaques, masses, or uniformly cover peritoneal surfaces. In most patients, death eventually results from locoregional progression within the abdominal cavity. In historical case series, treatment by palliative surgery, systemic or intraperitoneal chemotherapy, and abdominal irradiation has resulted in median survival of 12 months.

### **Treatment**

Surgical cytoreduction (resection of visible disease) in conjunction with hyperthermic intraperitoneal chemotherapy (HIPEC) is designed to remove visible tumor deposits and residual microscopic disease. By delivering chemotherapy intraperitoneally, drug exposure to the peritoneal surface is increased some 20-fold compared with systemic exposure. In addition, previous animal and in vitro studies have suggested that the cytotoxicity of mitomycin C is enhanced at temperatures greater than 39°C (102.2°F).

## Ovarian Cancer

Several different types of malignancies can arise in the ovaries; epithelial carcinoma is the most common, accounting for 90% of malignant ovarian tumors. Epithelial ovarian cancer is the fifth most common cause of cancer death in women in the United States. Most ovarian cancer patients (>70%) present with widespread disease, and annual mortality is 65% of the incidence rate. In addition, African American women reportedly have a higher prevalence of presenting with more advanced tumors, being undertreated or untreated, and having shorter disease-free survival compared to other racial groups.<sup>8</sup>

## Treatment

Current management of advanced epithelial ovarian cancer is cytoreductive surgery (CRS) followed by combination chemotherapy. Tumor recurrences are common, and the prognosis for recurrent disease is poor.

Cytoreductive surgery plus HIPEC in combination with systemic chemotherapy is being studied for primary and recurrent disease. Because HIPEC is administered at the time of surgery, treatment-related morbidity may be reduced compared with intraperitoneal chemotherapy administered postoperatively.

## Summary

### Description

Cytoreductive surgery (CRS) includes peritonectomy (ie, peritoneal stripping) procedures and multivisceral resections, depending on the extent of intra-abdominal tumor dissemination. Cytoreductive surgery may be followed by infusion of intraperitoneal chemotherapy with or without heating, which is intended to improve the tissue penetration of the chemotherapy. When heated, this is referred to as hyperthermic intraperitoneal chemotherapy (HIPEC). Cytoreductive surgery and HIPEC have been proposed for a number of intra-abdominal and pelvic malignancies such as pseudomyxoma peritonei and peritoneal carcinomatosis from colorectal, gastric, or endometrial cancer.

### Summary of Evidence

For individuals who have pseudomyxoma peritonei who receive cytoreductive surgery (CRS) plus hyperthermic intraperitoneal chemotherapy (HIPEC), the evidence includes cohort studies and a systematic review. Relevant outcomes are overall survival (OS), disease-specific survival, quality of life (QOL), and treatment-related mortality and morbidity. Retrospective cohort studies and systematic reviews have reported median survival ranging from 47 to 156 months and 5-year OS ranging from 41% to 96% for patients with primary treatment for pseudomyxoma peritonei treated with CRS plus HIPEC. Two retrospective studies reported results of CRS plus HIPEC for recurrence with 5-year OS rates of 34% and 79%. Although no direct comparisons between CRS plus HIPEC and other interventions have been published, traditional surgical debulking is not curative, and complete CRS alone (without HIPEC) has been associated with a 5-year OS of approximately 50%, along with high recurrence rates (91%, with a median disease-free survival of 24 months). Median progression-free survival (PFS) PFS with CRS plus HIPEC as primary treatment has been reported as 40 to 78 months, with 5-year PFS rates of 38% to 80%. Procedure-related morbidity and mortality have generally decreased over time. Because the prevalence of pseudomyxoma peritonei is very low, conducting comparative trials is difficult. The evidence is sufficient to determine that the technology results in an improvement in the net health outcome.

For individuals who have peritoneal carcinomatosis of colorectal origin who receive CRS plus HIPEC, the evidence includes randomized controlled trials (RCTs), systematic reviews, and a large number of observational studies. Relevant outcomes are OS, disease-specific survival, QOL, and treatment-related mortality and morbidity. Meta-analyses have found that CRS plus HIPEC, compared with traditional therapy without HIPEC, is associated with significantly higher survival rates and was not associated with significantly higher treatment-related morbidity rates. One RCT, in which patients with peritoneal carcinomatosis due to colorectal cancer were followed for at least 6 years, demonstrated improved survival in patients who received CRS plus HIPEC and systemic chemotherapy compared with patients who received systemic chemotherapy alone. However, procedure-related morbidity and mortality rates were relatively high, and systemic chemotherapy regimens did not use currently available biologic agents.

A more recent RCT found no survival benefit with CRS plus HIPEC over CRS alone, and a higher rate of adverse events 31 to 60 days post-procedure in the CRS plus HIPEC group. The lack of benefit seen with HIPEC in this trial may have been due to several factors, including the short duration of HIPEC treatment, the extensive use of preprocedural systemic chemotherapy, and the high rates of complete cytoreduction achieved in both groups. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

For individuals who have peritoneal carcinomatosis of gastric origin who receive CRS plus HIPEC, the evidence includes 2 small RCTs, observational studies, and 2 systematic reviews. Relevant outcomes are OS, disease-specific survival, QOL, and treatment-related mortality and morbidity. A 2017 meta-analysis identified 2 RCTs and 12 controlled nonrandomized studies comparing surgery plus HIPEC with standard surgical management in patients who had peritoneal carcinomatosis due to gastric cancer. One meta-analysis found significantly better survival in the surgery plus HIPEC group at 1 year but not at 2 or 3 years. Another meta-analysis found survival benefit was reported in the CRS plus HIPEC groups at 1, 2 and 3 years. One small (N=17) preliminary RCT showed improved survival in patients with peritoneal carcinomatosis due to gastric cancer who received CRS plus HIPEC compared with patients who received chemotherapy alone. Another (N=68) RCT showed improved survival in patients who received CRS plus HIPEC compared with CRS alone. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

For individuals who have peritoneal carcinomatosis of endometrial origin who receive CRS plus HIPEC, the evidence includes cohort studies. Relevant outcomes are OS, disease-specific survival, QOL, and treatment-related mortality and morbidity. Only uncontrolled retrospective cohort studies were available, with the largest including only 43 patients. Randomized trials that compare CRS plus HIPEC with standard treatment (eg, CRS alone or systemic chemotherapy alone) are needed. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

For individuals who have peritoneal mesothelioma who receive CRS plus HIPEC, the evidence includes retrospective cohort studies and systematic reviews. Relevant outcomes are OS, disease-specific survival, QOL, and treatment-related mortality and morbidity. Retrospective cohort studies have shown median and 5-year OS ranging from 30 to 92 months and from 33% to 68%, respectively, for patients with peritoneal mesothelioma treated with CRS plus HIPEC. Although no RCTs or comparative studies have been published, historical case series have reported a median survival of 12 months with treatment by palliative surgery, systemic or intraperitoneal chemotherapy, and abdominal irradiation. Procedure-related morbidity and mortality rates with CRS plus HIPEC have remained relatively steady over time, at approximately 35% and 5%, respectively. Because the prevalence of peritoneal mesothelioma is very low, conducting comparative trials is difficult. The evidence is sufficient to determine that the technology results in an improvement in the net health outcome.

For individuals who have newly diagnosed stage III ovarian cancer who receive CRS plus HIPEC, the evidence includes systematic reviews and RCTs ~~and~~ RCT. Relevant outcomes are OS, disease-specific survival, QOL, and treatment-related mortality and morbidity. For patients with newly diagnosed stage III ovarian cancer who had received neoadjuvant chemotherapy, HIPEC increased the time to disease recurrence and reduced mortality. It did not increase serious adverse events compared with surgery alone. The evidence is sufficient to determine that the technology results in an improvement in the net health outcome.

For individuals who have recurrent stage IIIC or IV ovarian cancer who receive CRS plus HIPEC, the evidence includes RCTs ~~and~~ RCT and systematic reviews. Relevant outcomes are OS, disease-specific survival, QOL, and treatment-related mortality and morbidity. For recurrent stage IIIC or IV disease (second-line setting), evidence from an RCT indicated that CRS plus HIPEC improved survival compared with CRS without HIPEC. However, interpretation of this study is limited because treatment groups in this RCT were unbalanced at baseline (variation in the completeness of cytoreduction), which has been shown to be associated with survival. Another RCT reported that CRS plus HIPEC did not result in superior outcomes compared to CRS without HIPEC for patients with platinum-sensitive recurrent

disease. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

For individuals who have appendiceal goblet cell tumors who receive CRS plus HIPEC, the evidence includes retrospective cohort studies. Relevant outcomes are OS, disease-specific survival, QOL, and treatment-related mortality and morbidity. A propensity score-matched analysis found that CRS plus HIPEC was associated with improved median survival compared to surgery alone. However, this analysis was limited by the retrospective nature of the data and small sample size (N=44). Rates of complete cytoreduction were not reported or accounted for in this study, so between-group differences in this or other variables may have influenced the observed outcomes. Additional studies- are needed. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

## Policy History

Date	Action
1/2024	Clarified coding information.
9/2023	Annual policy review. References added and updated. Policy statements unchanged.
9/2022	Annual policy review. References added and updated. Minor editorial refinements to policy statements; intent unchanged.
8/2021	Annual policy review. Description, summary, and references updated. Policy statements unchanged.
12/2020	Annual policy review. Description, summary, and references updated. Policy statements unchanged.
11/2019	Annual policy review. Description, summary, and references updated. Policy statements unchanged.
3/2019	Annual policy review. Policy statement updated to include hyperthermic intraperitoneal chemotherapy for the treatment of newly diagnosed stage III ovarian cancer. Policy title changed. Effective 3/1/2019.
1/2018	Clarified coding information.
8/2017	Annual policy review. New references added
8/2016	Annual policy review. New references added
6/2015	Annual policy review. New investigational indications described. Clarified coding information. Title changed to "Select Intra-Abdominal and Pelvic Malignancies." Effective 6/1/2015.
5/2014	Updated Coding section with ICD10 procedure and diagnosis codes. Effective 10/2015.
4/2014	Clarified coding information.
1/2014	Annual policy review. New references added.
11/2011-4/2012	Medical policy ICD10 remediation: Formatting, editing and coding updates. No changes to policy statements.
7/2011	Reviewed - Medical Policy Group – Hematology and Oncology. No changes to policy statements.
9/2010	Reviewed - Medical Policy Group - Hematology and Oncology. No changes to policy statements.
9/2009	Reviewed - Medical Policy Group - Hematology and Oncology. No changes to policy statements.
10/2008	Reviewed - Medical Policy Group – Hematology and Oncology. No changes to policy statements.
3/2008	Medical policy #048 describing covered and non-covered indications. Effective 3/1/2008.

## Information Pertaining to All Blue Cross Blue Shield Medical Policies

Click on any of the following terms to access the relevant information:



[Medical Policy Terms of Use](#)  
[Managed Care Guidelines](#)  
[Indemnity/PPO Guidelines](#)  
[Clinical Exception Process](#)  
[Medical Technology Assessment Guidelines](#)

## References

1. Maggiori L, Elias D. Curative treatment of colorectal peritoneal carcinomatosis: current status and future trends. *Eur J Surg Oncol.* Jul 2010; 36(7): 599-603. PMID 20605396
2. National Organization for Rare Disorders. Pseudomyxoma peritonei. <https://rarediseases.org/rare-diseases/pseudomyxoma-peritonei/>. Accessed May 18, 2023.
3. Elias D, Honoré C, Ciuchendéa R, et al. Peritoneal pseudomyxoma: results of a systematic policy of complete cytoreductive surgery and hyperthermic intraperitoneal chemotherapy. *Br J Surg.* Sep 2008; 95(9): 1164-71. PMID 18690633
4. Yonemura Y, Kawamura T, Bandou E, et al. Advances in the management of gastric cancer with peritoneal dissemination. *Recent Results Cancer Res.* 2007; 169: 157-64. PMID 17506258
5. Yonemura Y, Endou Y, Shinbo M, et al. Safety and efficacy of bidirectional chemotherapy for treatment of patients with peritoneal dissemination from gastric cancer: Selection for cytoreductive surgery. *J Surg Oncol.* Sep 15 2009; 100(4): 311-6. PMID 19697437
6. National Comprehensive Cancer Network (NCCN). NCCN Clinical practice guidelines in oncology: gastric cancer. Version 1.2023. [https://www.nccn.org/professionals/physician\\_gls/pdf/gastric.pdf](https://www.nccn.org/professionals/physician_gls/pdf/gastric.pdf). Accessed May 16, 2023.
7. Baratti D, Kusamura S, Deraco M. Diffuse malignant peritoneal mesothelioma: systematic review of clinical management and biological research. *J Surg Oncol.* Jun 2011; 103(8): 822-31. PMID 21283990
8. Chornokur G, Amankwah EK, Schildkraut JM, et al. Global ovarian cancer health disparities. *Gynecol Oncol.* Apr 2013; 129(1): 258-64. PMID 23266352
9. Glockzin G, Ghali N, Lang SA, et al. Results of cytoreductive surgery and hyperthermic intraperitoneal chemotherapy for peritoneal carcinomatosis from colorectal cancer. *J Surg Oncol.* Sep 15 2009; 100(4): 306-10. PMID 19697436
10. Jimenez W, Sardi A, Nieroda C, et al. Predictive and prognostic survival factors in peritoneal carcinomatosis from appendiceal cancer after cytoreductive surgery with hyperthermic intraperitoneal chemotherapy. *Ann Surg Oncol.* Dec 2014; 21(13): 4218-25. PMID 24986239
11. Glehen O, Gilly FN, Boutitie F, et al. Toward curative treatment of peritoneal carcinomatosis from nonovarian origin by cytoreductive surgery combined with perioperative intraperitoneal chemotherapy: a multi-institutional study of 1,290 patients. *Cancer.* Dec 15 2010; 116(24): 5608-18. PMID 20737573
12. Elias D, Gilly F, Quenet F, et al. Pseudomyxoma peritonei: a French multicentric study of 301 patients treated with cytoreductive surgery and intraperitoneal chemotherapy. *Eur J Surg Oncol.* May 2010; 36(5): 456-62. PMID 20227231
13. Chua TC, Yan TD, Smigielski ME, et al. Long-term survival in patients with pseudomyxoma peritonei treated with cytoreductive surgery and perioperative intraperitoneal chemotherapy: 10 years of experience from a single institution. *Ann Surg Oncol.* Jul 2009; 16(7): 1903-11. PMID 19387742
14. Vaira M, Cioppa T, DE Marco G, et al. Management of pseudomyxoma peritonei by cytoreduction+HIPEC (hyperthermic intraperitoneal chemotherapy): results analysis of a twelve-year experience. *In Vivo.* 2009; 23(4): 639-44. PMID 19567401
15. Marcotte E, Dubé P, Drolet P, et al. Hyperthermic intraperitoneal chemotherapy with oxaliplatin as treatment for peritoneal carcinomatosis arising from the appendix and pseudomyxoma peritonei: a survival analysis. *World J Surg Oncol.* Nov 07 2014; 12: 332. PMID 25380618
16. Yan TD, Black D, Savady R, et al. A systematic review on the efficacy of cytoreductive surgery and perioperative intraperitoneal chemotherapy for pseudomyxoma peritonei. *Ann Surg Oncol.* Feb 2007; 14(2): 484-92. PMID 17054002
17. Lord AC, Shihab O, Chandrakumaran K, et al. Recurrence and outcome after complete tumour removal and hyperthermic intraperitoneal chemotherapy in 512 patients with pseudomyxoma peritonei from perforated appendiceal mucinous tumours. *Eur J Surg Oncol.* Mar 2015; 41(3): 396-9. PMID 25216980
18. Sardi A, Jimenez WA, Nieroda C, et al. Repeated cytoreductive surgery and hyperthermic intraperitoneal chemotherapy in peritoneal carcinomatosis from appendiceal cancer: analysis of survival outcomes. *Eur J Surg Oncol.* Nov 2013; 39(11): 1207-13. PMID 24007834

19. Li J, Wang AR, Chen XD, et al. Effect of hyperthermic intraperitoneal chemotherapy in combination with cytoreductive surgery on the prognosis of patients with colorectal cancer peritoneal metastasis: a systematic review and meta-analysis. *World J Surg Oncol.* Jun 14 2022; 20(1): 200. PMID 35701802
20. Huang CQ, Min Y, Wang SY, et al. Cytoreductive surgery plus hyperthermic intraperitoneal chemotherapy improves survival for peritoneal carcinomatosis from colorectal cancer: a systematic review and meta-analysis of current evidence. *Oncotarget.* Aug 15 2017; 8(33): 55657-55683. PMID 28903452
21. Shan LL, Saxena A, Shan BL, et al. Quality of life after cytoreductive surgery and hyperthermic intraperitoneal chemotherapy for peritoneal carcinomatosis: A systematic review and meta-analysis. *Surg Oncol.* Dec 2014; 23(4): 199-210. PMID 25466850
22. Seretis C, Youssef H. Quality of life after cytoreductive surgery and intraoperative hyperthermic intraperitoneal chemotherapy for peritoneal surface malignancies: a systematic review. *Eur J Surg Oncol.* Dec 2014; 40(12): 1605-13. PMID 25242382
23. Quénet F, Elias D, Roca L, et al. Cytoreductive surgery plus hyperthermic intraperitoneal chemotherapy versus cytoreductive surgery alone for colorectal peritoneal metastases (PRODIGE 7): a multicentre, randomised, open-label, phase 3 trial. *Lancet Oncol.* Feb 2021; 22(2): 256-266. PMID 33476595
24. Verwaal VJ, van Ruth S, de Bree E, et al. Randomized trial of cytoreduction and hyperthermic intraperitoneal chemotherapy versus systemic chemotherapy and palliative surgery in patients with peritoneal carcinomatosis of colorectal cancer. *J Clin Oncol.* Oct 15 2003; 21(20): 3737-43. PMID 14551293
25. Verwaal VJ, Bruin S, Boot H, et al. 8-year follow-up of randomized trial: cytoreduction and hyperthermic intraperitoneal chemotherapy versus systemic chemotherapy in patients with peritoneal carcinomatosis of colorectal cancer. *Ann Surg Oncol.* Sep 2008; 15(9): 2426-32. PMID 18521686
26. Granieri S, Bonomi A, Frassini S, et al. Prognostic impact of cytoreductive surgery (CRS) with hyperthermic intraperitoneal chemotherapy (HIPEC) in gastric cancer patients: A meta-analysis of randomized controlled trials. *Eur J Surg Oncol.* Nov 2021; 47(11): 2757-2767. PMID 34001385
27. Desiderio J, Chao J, Melstrom L, et al. The 30-year experience-A meta-analysis of randomised and high-quality non-randomised studies of hyperthermic intraperitoneal chemotherapy in the treatment of gastric cancer. *Eur J Cancer.* Jul 2017; 79: 1-14. PMID 28456089
28. Rudloff U, Langan RC, Mullinax JE, et al. Impact of maximal cytoreductive surgery plus regional heated intraperitoneal chemotherapy (HIPEC) on outcome of patients with peritoneal carcinomatosis of gastric origin: results of the GYMSSA trial. *J Surg Oncol.* Sep 2014; 110(3): 275-84. PMID 25042700
29. Yang XJ, Huang CQ, Suo T, et al. Cytoreductive surgery and hyperthermic intraperitoneal chemotherapy improves survival of patients with peritoneal carcinomatosis from gastric cancer: final results of a phase III randomized clinical trial. *Ann Surg Oncol.* Jun 2011; 18(6): 1575-81. PMID 21431408
30. Navarro-Barrios Á, Gil-Martínez J, Ramos-Bernardo I, et al. Intraperitoneal hyperthermic chemotherapy after cytoreduction in patients with peritoneal metastases from endometrial cancer. The next frontier?. *Surg Oncol.* Jun 2020; 33: 19-23. PMID 32561085
31. Cornali T, Sammartino P, Kopanakis N, et al. Cytoreductive Surgery Plus Hyperthermic Intraperitoneal Chemotherapy for Patients with Peritoneal Metastases from Endometrial Cancer. *Ann Surg Oncol.* Mar 2018; 25(3): 679-687. PMID 29282600
32. Helm JH, Miura JT, Glenn JA, et al. Cytoreductive surgery and hyperthermic intraperitoneal chemotherapy for malignant peritoneal mesothelioma: a systematic review and meta-analysis. *Ann Surg Oncol.* May 2015; 22(5): 1686-93. PMID 25124472
33. Robella M, Vaira M, Mellano A, et al. Treatment of diffuse malignant peritoneal mesothelioma (DMPM) by cytoreductive surgery and HIPEC. *Minerva Chir.* Feb 2014; 69(1): 9-15. PMID 24675242
34. Alexander HR, Bartlett DL, Pingpank JF, et al. Treatment factors associated with long-term survival after cytoreductive surgery and regional chemotherapy for patients with malignant peritoneal mesothelioma. *Surgery.* Jun 2013; 153(6): 779-86. PMID 23489943
35. Yan TD, Deraco M, Baratti D, et al. Cytoreductive surgery and hyperthermic intraperitoneal chemotherapy for malignant peritoneal mesothelioma: multi-institutional experience. *J Clin Oncol.* Dec 20 2009; 27(36): 6237-42. PMID 19917862
36. Kim SI, Kim JH, Lee S, et al. Hyperthermic intraperitoneal chemotherapy for epithelial ovarian cancer: A meta-analysis. *Gynecol Oncol.* Dec 2022; 167(3): 547-556. PMID 36273925



37. Zhang G, Zhu Y, Liu C, et al. The prognosis impact of hyperthermic intraperitoneal chemotherapy (HIPEC) plus cytoreductive surgery (CRS) in advanced ovarian cancer: the meta-analysis. *J Ovarian Res.* Apr 17 2019; 12(1): 33. PMID 30995948
38. Wang Y, Ren F, Chen P, et al. Effects of CytoReductive surgery plus hyperthermic IntraPEritoneal chemotherapy (HIPEC) versus CytoReductive surgery for ovarian cancer patients: A systematic review and meta-analysis. *Eur J Surg Oncol.* Mar 2019; 45(3): 301-309. PMID 30786961
39. Antonio CCP, Alida GG, Elena GG, et al. Cytoreductive Surgery With or Without HIPEC After Neoadjuvant Chemotherapy in Ovarian Cancer: A Phase 3 Clinical Trial. *Ann Surg Oncol.* Apr 2022; 29(4): 2617-2625. PMID 34812982
40. van Driel WJ, Koole SN, Sikorska K, et al. Hyperthermic Intraperitoneal Chemotherapy in Ovarian Cancer. *N Engl J Med.* Jan 18 2018; 378(3): 230-240. PMID 29342393
41. Huo YR, Richards A, Liauw W, et al. Hyperthermic intraperitoneal chemotherapy (HIPEC) and cytoreductive surgery (CRS) in ovarian cancer: A systematic review and meta-analysis. *Eur J Surg Oncol.* Dec 2015; 41(12): 1578-89. PMID 26453145
42. Spiliotis J, Halkia E, Lianos E, et al. Cytoreductive surgery and HIPEC in recurrent epithelial ovarian cancer: a prospective randomized phase III study. *Ann Surg Oncol.* May 2015; 22(5): 1570-5. PMID 25391263
43. Zivanovic O, Chi DS, Zhou Q, et al. Secondary Cytoreduction and Carboplatin Hyperthermic Intraperitoneal Chemotherapy for Platinum-Sensitive Recurrent Ovarian Cancer: An MSK Team Ovary Phase II Study. *J Clin Oncol.* Aug 10 2021; 39(23): 2594-2604. PMID 34019431
44. Sluiter NR, van der Bilt JD, Croll DMR, et al. Cytoreduction and Hyperthermic Intraperitoneal Chemotherapy (HIPEC) Versus Surgery Without HIPEC for Goblet-Cell Carcinoids and Mixed Adenoneuroendocrine Carcinomas: Propensity Score-Matched Analysis of Centers in the Netherlands and Belgium. *Clin Colorectal Cancer.* Sep 2020; 19(3): e87-e99. PMID 32651131
45. McConnell YJ, Mack LA, Gui X, et al. Cytoreductive surgery with hyperthermic intraperitoneal chemotherapy: an emerging treatment option for advanced goblet cell tumors of the appendix. *Ann Surg Oncol.* Jun 2014; 21(6): 1975-82. PMID 24398544
46. Zambrano-Vera K, Sardi A, Munoz-Zuluaga C, et al. Outcomes in Peritoneal Carcinomatosis from Appendiceal Goblet Cell Carcinoma Treated with Cytoreductive Surgery and Hyperthermic Intraperitoneal Chemotherapy (CRS/HIPEC). *Ann Surg Oncol.* Jan 2020; 27(1): 179-187. PMID 31646450
47. Morris VK, Kennedy EB, Baxter NN, et al. Treatment of Metastatic Colorectal Cancer: ASCO Guideline. *J Clin Oncol.* Jan 20 2023; 41(3): 678-700. PMID 36252154
48. Vogel JD, Felder SI, Bhama AR, et al. The American Society of Colon and Rectal Surgeons Clinical Practice Guidelines for the Management of Colon Cancer. *Dis Colon Rectum.* Feb 01 2022; 65(2): 148-177. PMID 34775402
49. Glasgow SC, Gaertner W, Stewart D, et al. The American Society of Colon and Rectal Surgeons, Clinical Practice Guidelines for the Management of Appendiceal Neoplasms. *Dis Colon Rectum.* Dec 2019; 62(12): 1425-1438. PMID 31725580
50. National Comprehensive Cancer Network (NCCN). NCCN Clinical practice guidelines in oncology: colon cancer. Version 2.2023. [https://www.nccn.org/professionals/physician\\_gls/pdf/colon.pdf](https://www.nccn.org/professionals/physician_gls/pdf/colon.pdf). Accessed May 17, 2023.
51. National Comprehensive Cancer Network (NCCN). NCCN Clinical practice guidelines in oncology: uterine neoplasms. Version 2.2023. [https://www.nccn.org/professionals/physician\\_gls/pdf/uterine.pdf](https://www.nccn.org/professionals/physician_gls/pdf/uterine.pdf). Accessed May 18, 2023.
52. National Comprehensive Cancer Network (NCCN). NCCN Clinical practice guidelines in oncology: rectal cancer. Version 2.2023. [https://www.nccn.org/professionals/physician\\_gls/pdf/rectal.pdf](https://www.nccn.org/professionals/physician_gls/pdf/rectal.pdf). Accessed May 15, 2023.
53. National Comprehensive Cancer Network (NCCN). NCCN Clinical practice guidelines in oncology: ovarian cancer including fallopian tube cancer and primary peritoneal cancer. Version 1.2023. [https://www.nccn.org/professionals/physician\\_gls/pdf/ovarian.pdf](https://www.nccn.org/professionals/physician_gls/pdf/ovarian.pdf). Accessed May 19, 2023.
54. Hoppenot C, Schuitevoerder D, Izquierdo FJ, et al. The Chicago Consensus on Peritoneal Surface Malignancies: Management of Ovarian Neoplasms. *Ann Surg Oncol.* Jun 2020; 27(6): 1780-1787. PMID 32285271

55. Izquierdo FJ, Schuitevoerder D, Plana A, et al. The Chicago Consensus on Peritoneal Surface Malignancies: Management of Colorectal Metastases. *Ann Surg Oncol.* Jun 2020; 27(6): 1761-1767. PMID 32285270
56. Izquierdo FJ, Schuitevoerder D, Plana A, et al. The Chicago Consensus on Peritoneal Surface Malignancies: Management of Gastric Metastases. *Ann Surg Oncol.* Jun 2020; 27(6): 1768-1773. PMID 32285269
57. Schuitevoerder D, Izquierdo FJ, Plana A, et al. The Chicago Consensus on Peritoneal Surface Malignancies: Management of Peritoneal Mesothelioma. *Ann Surg Oncol.* Jun 2020; 27(6): 1774-1779. PMID 32285273
58. Schuitevoerder D, Plana A, Izquierdo FJ, et al. The Chicago Consensus on Peritoneal Surface Malignancies: Management of Appendiceal Neoplasms. *Ann Surg Oncol.* Jun 2020; 27(6): 1753-1760. PMID 32285275