



# MASSACHUSETTS

Blue Cross Blue Shield of Massachusetts is an Independent Licensee of the Blue Cross and Blue Shield Association

## Medical Policy

# Radiofrequency Coblation Tenotomy for Musculoskeletal Conditions

### Table of Contents

- [Policy: Commercial](#)
- [Coding Information](#)
- [Information Pertaining to All Policies](#)
- [Policy: Medicare](#)
- [Description](#)
- [References](#)
- [Authorization Information](#)
- [Policy History](#)

### Policy Number: 080

BCBSA Reference Number: 7.01.165 (For Plans internal use only)

NCD/LCD:

### Related Policies

- Diagnosis and Treatment of Sacroiliac Joint Pain, [#320](#)
- Decompression of the Intervertebral Disc Using Laser Energy (Laser Discectomy) or Radiofrequency-Coblation (Nucleoplasty), [#271](#)

### Policy

## Commercial Members: Managed Care (HMO and POS), PPO, and Indemnity Medicare HMO Blue<sup>SM</sup> and Medicare PPO Blue<sup>SM</sup> Members

Radiofrequency coblation tenotomy is considered **INVESTIGATIONAL** as a treatment for musculoskeletal conditions, including but not limited to the following conditions:

- Plantar fasciitis
- Lateral epicondylitis
- Wrist tendinopathy
- Shoulder or rotator cuff tendinopathy
- Achilles tendinopathy
- Patellar tendinopathy.

### Prior Authorization Information

#### Inpatient

- For services described in this policy, precertification/preauthorization **IS REQUIRED** for all products if the procedure is performed **inpatient**.

#### Outpatient

- For services described in this policy, see below for products where prior authorization **might be required** if the procedure is performed **outpatient**.

	Outpatient
Commercial Managed Care (HMO and POS)	This is <b>not</b> a covered service.

<b>Commercial PPO and Indemnity</b>	This is <b>not</b> a covered service.
<b>Medicare HMO Blue<sup>SM</sup></b>	This is <b>not</b> a covered service.
<b>Medicare PPO Blue<sup>SM</sup></b>	This is <b>not</b> a covered service.

## CPT Codes / HCPCS Codes / ICD Codes

*Inclusion or exclusion of a code does not constitute or imply member coverage or provider reimbursement. Please refer to the member's contract benefits in effect at the time of service to determine coverage or non-coverage as it applies to an individual member.*

*Providers should report all services using the most up-to-date industry-standard procedure, revenue, and diagnosis codes, including modifiers where applicable.*

There is no specific CPT code for this procedure.

## Description

### Radiofrequency Coblation

Radiofrequency (RF) coblation uses bipolar low-frequency energy in an electrically conductive fluid (eg, saline) to generate a high-density plasma field around the energy source. This creates a low-temperature field of ionizing particles that break organic bonds within the target tissue. Coblation technology is used in a variety of surgical procedures, particularly related to otolaryngology and orthopedics. The proposed advantage of coblation is that the procedure provides for controlled and highly localized ablation, resulting in minimal damage to surrounding tissue. Radiofrequency coblation was also found to exhibit several properties that may make it an attractive option for addressing the underlying pathophysiology of chronic tendinopathies, namely increased angiogenesis, reduction of inflammatory responses, and increased expression of growth factors.<sup>1</sup> Radiofrequency coblation surgical wands are utilized by orthopedic surgeons in minimally invasive arthroscopic procedures to facilitate soft tissue debridement, subacromial decompression, meniscal removal and sculpting, or tendon debridement.

The use of coblation technology for disc nucleoplasty and sacroiliac joint pain is addressed separately in policy #[320](#) and policy #[271](#).

### Tendinopathy

Tendinopathy is a clinical pain syndrome characterized by tendon thickening due to proliferation and chronic irritation of neovascular repair tissue with a history of repetitive tendon loading. This condition commonly results from overuse and has a high incidence rate in athletes and laborers. Clinical history should clarify predisposing training or activity and assess the level of functioning. Biomechanical abnormalities during activity should be identified and corrected. Standard treatment may, therefore, consist of biomechanical modification, activity modification, physical therapy (eg, heavy load resistance training), and nonsteroidal anti-inflammatory medication. For chronic tendinopathies, glucocorticoids should only be used in select cases (eg, rotator cuff tendinopathy). Surgical consultation following 6 months of a well-designed physical therapy program with adjunct medical treatments can be considered if there is no improvement in pain or function.<sup>2</sup> Validated and reliable functional assessment scores should be utilized by the clinician to grade symptoms and assess patient function. Examples of suitable scales include the Victoria Institute of Sport Assessment for Achilles tendinopathy.<sup>3</sup> Surgical approaches may involve incisions to the paratendon and removal of adhesions and degenerate tissue. Longitudinal incisions may be made in the tendon to promote a repair response. This latter strategy has also been delivered via minimally invasive arthroscopic approaches.<sup>4,5</sup> These approaches may also address the debridement of the neovascular supply to the tendon surface. Collectively, a prolonged recovery duration to accommodate tendon healing may be required with these interventions.

### Plantar Fasciitis

Plantar fasciitis is a musculoskeletal condition characterized by pain in the plantar region of the foot that worsens upon initiation of walking and with local point tenderness elicited during a clinical examination. Radiographic and ultrasonographic studies are not typically indicated for primary diagnosis but may be useful in ruling out alternative causes and visualizing the thickening of the plantar fascia. Initial standard

therapy may consist of stretching exercises, orthotics, activity and lifestyle modification, nonsteroidal anti-inflammatory drugs, splints or casts, and glucocorticoid injections. The vast majority of patients improve without surgery. Surgery is generally considered a last line of therapy and is reserved for individuals who do not respond to at least 6 to 12 months of initial, nonsurgical therapy. Surgical approaches include variations of open or endoscopic, partial or complete, plantar fascia release, which may or may not include calcaneal spur resection, excision of abnormal tissue, and nerve decompression. The use of RF microtenotomy during open or percutaneous surgery has been explored alone or in combination with plantar fasciotomy.<sup>6</sup>

Plantar fasciitis is one of the most common causes of foot and heel pain in adults. It is estimated to be responsible for approximately 1 million patient medical visits per year in the U.S.<sup>7</sup> The peak incidence of the condition in the general population occurs between ages 40 and 60. There is a higher incidence rate among runners with a younger age of onset. The etiology of plantar fasciitis is poorly understood and may be multifactorial in nature. Contributing risk factors may include obesity, prolonged standing or activity, flat feet, and reduced ankle dorsiflexion.<sup>8,9</sup> Plantar fasciitis has been reported in association with fluoride use for the treatment of osteoporosis.<sup>10</sup> Differential sources of foot and heel pain may include Achilles tendinopathy, stress fractures due to osteoporosis, rheumatoid arthritis, peripheral neuropathies associated with diabetes, extrinsic factors (eg, inappropriate footwear), aging, and structural disorders.

### **Lateral Epicondylitis**

Lateral epicondylitis, also known as tennis elbow, represents chronic tendinosis of the myotendinous group of the lateral epicondyle characterized by pain and disability. The incidence in the general population may approach 1% to 3%.<sup>11</sup> Risk factors include smoking, obesity, forceful activity, and repetitive activity for at least 2 hours daily. Lateral epicondylitis is characterized by injury to the extensor carpi radialis brevis or extensor digitorum communis muscles. The condition is diagnosed through findings of localized tenderness and pain with clinical examination. Initial conservative management includes modification of activity and biomechanics, counterforce bracing or splinting, nonsteroidal anti-inflammatory drugs, and physical therapy.<sup>12</sup> Surgical referral is typically reserved for patients with severe symptoms that do not improve despite compliance with an appropriately designed physical therapy program for at least 6 months.

## **Summary**

Radiofrequency (RF) coblation is being evaluated for the treatment of plantar fasciitis, lateral epicondylitis, and various musculoskeletal tendinopathies. When utilized for tenotomy, bipolar RF energy is directed into the tendon to generate a controlled, low-temperature field of ionizing particles that break organic bonds, ablating or debriding target tissue with the goal of relieving pain and restoring function.

### **Summary of Evidence**

For individuals with plantar fasciitis who receive RF coblation tenotomy, the evidence includes nonrandomized, comparative cohort studies and case series. Relevant outcomes are symptoms, functional outcomes, QOL, medication use, and treatment-related morbidity. The trials reported improved pain and functional scores over 3 to 12 months, with improved outcomes with open versus percutaneous approaches. However, open RF coblation microtenotomy was associated with a higher incidence of postoperative persistent pain (9.1%) compared to endoscopic plantar fasciotomy (0%) in 1 study, with a separate study reporting a complication rate of 33% when both interventions were used in combination. A higher number of postoperative pain recurrences at 6 and 12 months were also reported with open RF coblation microtenotomy compared to endoscopic plantar fasciotomy. The durability of this intervention is unknown as no studies have reported long-term outcomes beyond 12 months. Studies are limited by small sample sizes, heterogeneity in surgical technique (open, percutaneous, endoscopic), missing data and/or inappropriate exclusions, lack of randomization, unclear blinding practices for patient outcome assessments, and poor statistical reporting. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

For individuals with lateral epicondylitis who receive RF coblation tenotomy, the evidence includes small RCTs. Relevant outcomes are symptoms, functional outcomes, QOL, medication use, and treatment-related morbidity. The trials compared RF microtenotomy to open or arthroscopic elbow release surgery.

Clinically meaningful improvements in pain and functional scores were noted for all treatment arms, with no significant differences between groups through 1 to 7 years of follow-up. For disability assessments in 1 study, open release surgery met the threshold for a clinically meaningful improvement over RF microtenotomy at 1 year, though this mean difference was not statistically significant. Studies were generally underpowered or demonstrated inconsistent delivery and unclear blinding of outcome assessments and inappropriate handling of missing or crossover data. No studies featuring RF coblation tenotomy for the treatment of wrist tendinopathy were identified. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

For individuals with Achilles tendinopathy who receive RF coblation tenotomy, the evidence includes small, single-blind RCTs. Relevant outcomes are symptoms, functional outcomes, QOL, medication use, and treatment-related morbidity. One trial did not demonstrate an added benefit for RF microdebridement compared to surgical decompression. Pain and functional outcomes improved in both groups but were not statistically different at a 6 month follow-up. The study was limited by a control group that showed significantly less severe symptom scores at baseline that did not fully meet the 2 point threshold for a clinically meaningful difference in pain score reduction. The other small RCT demonstrated potential benefits in pain and quality of life for RF microtenotomy (ArthroCare) compared with physical therapy at 2 years. But, conclusions cannot be drawn based on these findings due to numerous notable study limitations. Larger, adequately controlled studies with longer follow-up durations are lacking. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

For individuals with shoulder or rotator cuff tendinopathy who receive RF coblation tenotomy, the evidence includes small RCTs. Relevant outcomes are symptoms, functional outcomes, QOL, medication use, and treatment-related morbidity. Trials did not demonstrate an added benefit for RF microdebridement compared to arthroscopic subacromial decompression surgery. Pain and functional outcomes improved in both groups but were not statistically different through 1 to 2 years of follow-up. Neither study prespecified a clinically meaningful difference in outcome measures nor were harms assessed throughout their course. The loss to follow-up in 1 study was 18.7%. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

For individuals with patellar tendinopathy who receive RF coblation tenotomy, the evidence includes 1 small RCT. Relevant outcomes are symptoms, functional outcomes, QOL, medication use, and treatment-related morbidity. The trial did not demonstrate an added benefit for RF microdebridement compared to mechanical debridement in patients with chondral lesions and patellar tendinopathy. The study lacked reporting with validated pain measures over time and reported a higher incidence of crepitus in patients undergoing RF microdebridement. Furthermore, the study only enrolled female participants, limiting the broader applicability of these findings. Larger studies with validated pain and functional outcome measures are required to adequately assess the technology. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

## Policy History

Date	Action
2/2024	Annual policy review. Policy updated with literature review through November 3, 2023; references added. Policy statements unchanged.
2/2023	Annual policy review. Description, summary, and references updated. Policy statements unchanged.
2/2022	Annual policy review. Description, summary, and references updated. Policy statements unchanged.
5/2020	New medical policy describing investigational indications. Effective 5/1/2020.

## Information Pertaining to All Blue Cross Blue Shield Medical Policies

Click on any of the following terms to access the relevant information:

[Medical Policy Terms of Use](#)

[Managed Care Guidelines](#)

## References

1. Tay KS, Ng YC, Singh IR, et al. Open technique is more effective than percutaneous technique for TOPAZ radiofrequency coblation for plantar fasciitis. *Foot Ankle Surg.* Dec 2012; 18(4): 287-92. PMID 23093126
2. Davenport TE, Kulig K, Matharu Y, et al. The EdUReP model for nonsurgical management of tendinopathy. *Phys Ther.* Oct 2005; 85(10): 1093-103. PMID 16180958
3. Robinson JM, Cook JL, Purdam C, et al. The VISA-A questionnaire: a valid and reliable index of the clinical severity of Achilles tendinopathy. *Br J Sports Med.* Oct 2001; 35(5): 335-41. PMID 11579069
4. Lohrer H, David S, Nauck T. Surgical treatment for achilles tendinopathy - a systematic review. *BMC Musculoskelet Disord.* May 10 2016; 17: 207. PMID 27165287
5. Nirschl RP, Pettrone FA. Tennis elbow. The surgical treatment of lateral epicondylitis. *J Bone Joint Surg Am.* Sep 1979; 61(6A): 832-9. PMID 479229
6. Chou AC, Ng SY, Su DH, et al. Radiofrequency microtenotomy is as effective as plantar fasciotomy in the treatment of recalcitrant plantar fasciitis. *Foot Ankle Surg.* Dec 2016; 22(4): 270-273. PMID 27810027
7. Riddle DL, Schappert SM. Volume of ambulatory care visits and patterns of care for patients diagnosed with plantar fasciitis: a national study of medical doctors. *Foot Ankle Int.* May 2004; 25(5): 303-10. PMID 15134610
8. Riddle DL, Pulisic M, Pidcoke P, et al. Risk factors for Plantar fasciitis: a matched case-control study. *J Bone Joint Surg Am.* May 2003; 85(5): 872-7. PMID 12728038
9. Rano JA, Fallat LM, Savoy-Moore RT. Correlation of heel pain with body mass index and other characteristics of heel pain. *J Foot Ankle Surg.* 2001; 40(6): 351-6. PMID 11777230
10. Riggs BL, Hodgson SF, Hoffman DL, et al. Treatment of primary osteoporosis with fluoride and calcium. Clinical tolerance and fracture occurrence. *JAMA.* Feb 01 1980; 243(5): 446-9. PMID 7351765
11. Shiri R, Viikari-Juntura E, Varonen H, et al. Prevalence and determinants of lateral and medial epicondylitis: a population study. *Am J Epidemiol.* Dec 01 2006; 164(11): 1065-74. PMID 16968862
12. Struijs PA, Kerkhoffs GM, Assendelft WJ, et al. Conservative treatment of lateral epicondylitis: brace versus physical therapy or a combination of both-a randomized clinical trial. *Am J Sports Med.* Mar 2004; 32(2): 462-9. PMID 14977675
13. Hamlin K, Munro C, Barker SL, et al. Open release versus radiofrequency microtenotomy in the treatment of lateral epicondylitis: a prospective randomized controlled trial. *Shoulder Elbow.* Jan 2018; 10(1): 45-51. PMID 29276537
14. Wang W, Rikhray IS, Chou ACC, et al. Endoscopic Plantar Fasciotomy vs Open Radiofrequency Microtenotomy for Recalcitrant Plantar Fasciitis. *Foot Ankle Int.* Jan 2018; 39(1): 11-17. PMID 29182482
15. Nayar SK, Alcock H, Vemulapalli K. Surgical treatment options for plantar fasciitis and their effectiveness: a systematic review and network meta-analysis. *Arch Orthop Trauma Surg.* Aug 2023; 143(8): 4641-4651. PMID 36596990
16. Yuan Y, Qian Y, Lu H, et al. Comparison of the therapeutic outcomes between open plantar fascia release and percutaneous radiofrequency ablation in the treatment of intractable plantar fasciitis. *J Orthop Surg Res.* Feb 18 2020; 15(1): 55. PMID 32070392
17. Huang DM, Chou AC, Yeo NE, et al. Radiofrequency Microtenotomy with Concurrent Gastrocnemius Recession Improves Postoperative Vitality Scores in the Treatment of Recalcitrant Plantar Fasciitis. *Ann Acad Med Singap.* Dec 2018; 47(12): 509-515. PMID 30636267
18. Weil L, Glover JP, Weil LS. A new minimally invasive technique for treating plantar fasciosis using bipolar radiofrequency: a prospective analysis. *Foot Ankle Spec.* Feb 2008; 1(1): 13-8. PMID 19825686
19. Yeap EJ, Chong KW, Yeo W, et al. Radiofrequency coblation for chronic foot and ankle tendinosis. *J Orthop Surg (Hong Kong).* Dec 2009; 17(3): 325-30. PMID 20065374
20. Sean NY, Singh I, Wai CK. Radiofrequency microtenotomy for the treatment of plantar fasciitis shows good early results. *Foot Ankle Surg.* Dec 2010; 16(4): 174-7. PMID 21047605

21. Meknas K, Al Hassoni TN, Odden-Miland Å, et al. Medium-Term Results After Treatment of Recalcitrant Lateral Epicondylitis: A Prospective, Randomized Study Comparing Open Release and Radiofrequency Microtenotomy. *Orthop J Sports Med.* Sep 2013; 1(4): 2325967113505433. PMID 26535247
22. Lee JH, Park I, Hyun HS, et al. A Comparison of Radiofrequency-Based Microtenotomy and Arthroscopic Release of the Extensor Carpi Radialis Brevis Tendon in Recalcitrant Lateral Epicondylitis: A Prospective Randomized Controlled Study. *Arthroscopy.* May 2018; 34(5): 1439-1446. PMID 29366739
23. Morrison RJM, Brock TM, Reed MR, et al. Radiofrequency Microdebridement Versus Surgical Decompression for Achilles Tendinosis: A Randomized Controlled Trial. *J Foot Ankle Surg.* 2017; 56(4): 708-712. PMID 28495412
24. Martin RL, Chimenti R, Cuddeford T, et al. Achilles Pain, Stiffness, and Muscle Power Deficits: Midportion Achilles Tendinopathy Revision 2018. *J Orthop Sports Phys Ther.* May 2018; 48(5): A1-A38. PMID 29712543
25. Al-Ani Z, Meknas D, Kartus JT, et al. Radiofrequency Microtenotomy or Physical Therapy for Achilles Tendinopathy: Results of a Randomized Clinical Trial. *Orthop J Sports Med.* Dec 2021; 9(12): 23259671211062555. PMID 34988234
26. Shibuya N, Thorud JC, Humphers JM, et al. Is percutaneous radiofrequency coblation for treatment of Achilles tendinosis safe and effective?. *J Foot Ankle Surg.* 2012; 51(6): 767-71. PMID 22974813
27. Lu Y, Zhang Q, Zhu Y, et al. Is radiofrequency treatment effective for shoulder impingement syndrome? A prospective randomized controlled study. *J Shoulder Elbow Surg.* Nov 2013; 22(11): 1488-94. PMID 23994459
28. Al-Ani Z, Jacobsen EW, Kartus JT, et al. Radiofrequency microtenotomy: a promising method for treatment of rotator cuff tendinopathy. *Knee Surg Sports Traumatol Arthrosc.* Dec 2019; 27(12): 3856-3863. PMID 31473769
29. Owens BD, Stickles BJ, Balikian P, et al. Prospective analysis of radiofrequency versus mechanical debridement of isolated patellar chondral lesions. *Arthroscopy.* Feb 2002; 18(2): 151-5. PMID 11830808
30. Schneider HP, Baca JM, Carpenter BB, et al. American College of Foot and Ankle Surgeons Clinical Consensus Statement: Diagnosis and Treatment of Adult Acquired Infracalcaneal Heel Pain. *J Foot Ankle Surg.* 2018; 57(2): 370-381. PMID 29284574
31. Hegmann KT, Hoffman HE, Belcourt RM, et al. ACOEM practice guidelines: elbow disorders. *J Occup Environ Med.* Nov 2013; 55(11): 1365-74. PMID 23963225
32. Centers for Medicare & Medicaid Services (CMS). National Coverage Determination for Thermal Intradiscal Procedures (TIPs) (150.11). 2009; <https://www.cms.gov/medicare-coverage-database/details/ncd-details.aspx?NCDId=324>. Accessed November 21, 2022.