



MASSACHUSETTS

Blue Cross Blue Shield of Massachusetts is an Independent Licensee of the Blue Cross and Blue Shield Association

Medical Policy

Bioimpedance Devices for the Detection of Lymphedema

Table of Contents

- [Policy: Commercial](#)
- [Policy: Medicare](#)
- [Authorization Information](#)
- [Coding Information](#)
- [Description](#)
- [Policy History](#)
- [Information Pertaining to All Policies](#)
- [References](#)

Policy Number: 261

BCBSA Reference Number: 2.01.82 (For Plan internal use only)

NCD/LCD: N/A

Related Policies

- Pneumatic Compression Pumps for Treatment of Lymphedema, #[354](#)
- Surgical and Debulking Treatments for Lymphedema, #[037](#)

Policy

Commercial Members: Managed Care (HMO and POS), PPO, and Indemnity Medicare HMO BlueSM and Medicare PPO BlueSM Members

Devices using bioimpedance (bioelectrical impedance spectroscopy) are considered [INVESTIGATIONAL](#) for use in the diagnosis, surveillance, or treatment of individuals with lymphedema, including use in subclinical secondary lymphedema.

Prior Authorization Information

Inpatient

- For services described in this policy, precertification/preauthorization **IS REQUIRED** for all products if the procedure is performed **inpatient**.

Outpatient

- For services described in this policy, see below for products where prior authorization **might be required** if the procedure is performed **outpatient**.

	Outpatient
Commercial Managed Care (HMO and POS)	This is not a covered service.
Commercial PPO and Indemnity	This is not a covered service.
Medicare HMO Blue SM	This is not a covered service.
Medicare PPO Blue SM	This is not a covered service.

CPT Codes / HCPCS Codes / ICD Codes

Inclusion or exclusion of a code does not constitute or imply member coverage or provider reimbursement. Please refer to the member's contract benefits in effect at the time of service to determine coverage or non-coverage as it applies to an individual member.

Providers should report all services using the most up-to-date industry-standard procedure, revenue, and diagnosis codes, including modifiers where applicable.

The following codes are included below for informational purposes only; this is not an all-inclusive list.

The following CPT code is considered investigational for Commercial Members: Managed Care (HMO and POS), PPO, Indemnity, Medicare HMO Blue and Medicare PPO Blue:

CPT Codes

CPT codes:	Code Description
93702	Bioimpedance spectroscopy (BIS), extracellular fluid analysis for lymphedema assessment(s)

Description

Lymphedema

Lymphedema is an accumulation of fluid due to disruption of lymphatic drainage. Lymphedema can be caused by congenital or inherited abnormalities in the lymphatic system (primary lymphedema) but is most often caused by acquired damage to the lymphatic system (secondary lymphedema). Breast cancer treatment is one of the most common causes of secondary lymphedema. Both the surgical removal of lymph nodes and radiotherapy are associated with development lymphedema in patients with breast cancer. In a systematic review of 72 studies (N=29612 women), DiSipio et al (2013) reported that approximately 1 in 5 women who survive breast cancer will develop arm lymphedema.¹ Risk factors for development of lymphedema that had a strong level of evidence were extensive surgery (ie, axillary-lymph-node dissection, greater number of lymph nodes dissected, mastectomy) and being overweight or obese.

Diagnosis and Staging

A diagnosis of secondary lymphedema is based on history (e.g., cancer treatment, trauma) and physical examination (localized, progressive edema and asymmetric limb measurements) when other causes of edema can be excluded. Imaging, such as MRI, computed tomography, ultrasound, or lymphoscintigraphy, may be used to differentiate lymphedema from other causes of edema in diagnostically challenging cases. Table 1 lists International Society of Lymphology guidance for staging lymphedema based on "softness" or "firmness" of the limb and the changes with an elevation of the limb.²

Table 1. Recommendations for Staging Lymphedema

Stage	Description
Stage 0 (subclinical)	Swelling is not evident, and most patients are asymptomatic despite impaired lymphatic transport
Stage I (mild)	Accumulation of fluid that subsides with limb elevation; soft edema that may pit, without evidence of dermal fibrosis
Stage II (moderate)	Does not resolve with limb elevation alone; limb may no longer pit on examination
Stage III (severe)	Lymphostatic elephantiasis; pitting can be absent; skin has trophic changes

Management and Treatment

Lymphedema is treated using elevation, compression, and exercise. Conservative therapy may consist of several features depending on the severity of the lymphedema. Patients are educated on the importance of self-care including hygiene practices to prevent infection, maintaining ideal body weight through diet and exercise, and limb elevation. Compression therapy consists of repeatedly applying padding and bandages or compression garments. Manual lymphatic drainage is a light pressure massage performed by trained physical therapists or by patients designed to move fluid from obstructed areas into functioning lymph

vessels and lymph nodes. Complete decongestive therapy is a multiphase treatment program involving all of the previously mentioned conservative treatment components at different intensities. Pneumatic compression pumps may also be considered as an adjunct to conservative therapy or as an alternative to self-manual lymphatic drainage in patients who have difficulty performing self-manual lymphatic drainage. In patients with more advanced lymphedema after fat deposition and tissue fibrosis has occurred, palliative surgery using reductive techniques such as liposuction may be performed.

Bioimpedance Spectroscopy

Bioimpedance spectroscopy is based on the theory that the level of opposition to the flow of electric current (impedance) through the body is inversely proportional to the volume of fluid in the tissue. In lymphedema, with the accumulation of excess interstitial fluid, tissue impedance decreases.

Bioimpedance has been proposed as a diagnostic test for this condition. In usual care, lymphedema is recognized clinically or via limb measurements. However, management via bioelectrical impedance spectroscopy has been proposed as a way to implement early treatment of subclinical lymphedema to potentially reduce its severity.

Summary

Description

Secondary lymphedema may develop following treatment for breast cancer. Bioimpedance, which uses resistance to electrical current to compare the composition of fluid compartments, could be used as a tool to diagnose lymphedema.

Summary of Evidence

For individuals who have known or suspected lymphedema who receive bioimpedance spectroscopy (BIS), the evidence includes systematic reviews, 1 randomized controlled trial (RCT), 1 prospective comparative observational study, and multiple uncontrolled observational studies. Relevant outcomes are test validity, symptoms, and quality of life. Diagnostic accuracy studies have found a poor correlation between bioimpedance analysis and the reference standard (volume displacement or circumferential measurement). Results from the PREVENT RCT comparing bioimpedance with standard tape measure following treatment for breast cancer have been published. At a median follow-up of 32.9 months, BIS patients triggered intervention at a lower rate than tape measured patients (20.1% vs 27.5%) and fewer patients progressed in this group (7.9% vs 19.2%). The RCT was limited by its open-label design and lack of reporting of important health outcomes. The single prospective comparative study found a significantly lower rate of clinical lymphedema in patients managed with BIS devices but had several limitations, including nonrandomized design, lack of blinding, lack of complete data on a substantial proportion of enrolled patients, and lack of a systematic method for diagnosing lymphedema in the control group. Retrospective studies suggested that postoperative bioimpedance monitoring is feasible but provide limited information about its efficacy. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

Policy History

Date	Action
3/2024	Annual policy review. Policy updated with literature review through November 27, 2023; references added. Policy statement unchanged.
3/2023	Annual policy review. Minor editorial refinements to policy statements; intent unchanged.
2/2022	Annual policy review. Description, summary, and references updated. Policy statements unchanged.
3/2021	Annual policy review. Description, summary, and references updated. Policy statements unchanged.
3/2020	Annual policy review. Description, summary, and references updated. Policy statements unchanged.
3/2019	Annual policy review. Description, summary, and references updated. Policy statements unchanged.

7/2017	Annual policy review. New references added.
3/2016	Annual policy review. New references added.
1/2015	Clarified coding information.
12/2014	Annual policy review. New references added.
2/2014	Annual policy review. New references added.
11/2011-4/2012	Medical policy ICD 10 remediation: Formatting, editing and coding updates. No changes to policy statements.
7/2011	Reviewed - Medical Policy Group – Hematology and Oncology. No changes to policy statements.
9/29/2010	Medical Policy 261 created.

Information Pertaining to All Blue Cross Blue Shield Medical Policies

Click on any of the following terms to access the relevant information:

[Medical Policy Terms of Use](#)

[Managed Care Guidelines](#)

[Indemnity/PPO Guidelines](#)

[Clinical Exception Process](#)

[Medical Technology Assessment Guidelines](#)

References

1. DiSipio T, Rye S, Newman B, et al. Incidence of unilateral arm lymphoedema after breast cancer: a systematic review and meta-analysis. *Lancet Oncol.* May 2013; 14(6): 500-15. PMID 23540561
2. International Society of Lymphology Executive Committee. The Diagnosis and Treatment of Peripheral Lymphedema: 2020 Consensus Document of the International Society of Lymphology. 2020; <https://isl.arizona.edu/sites/default/files/2021-09/Consensus%20Document-SM.pdf>. Accessed November 27, 2023.
3. Pusic AL, Cemal Y, Albornoz C, et al. Quality of life among breast cancer patients with lymphedema: a systematic review of patient-reported outcome instruments and outcomes. *J Cancer Surviv.* Mar 2013; 7(1): 83-92. PMID 23212603
4. Oremus M, Walker K, Dayes I, et al. Technology Assessment: Diagnosis and treatment of secondary lymphedema. Rockville, MD: Agency for Healthcare Research and Quality; 2010.
5. Cornish BH, Chapman M, Hirst C, et al. Early diagnosis of lymphedema using multiple frequency bioimpedance. *Lymphology.* Mar 2001; 34(1): 2-11. PMID 11307661
6. Hayes S, Janda M, Cornish B, et al. Lymphedema secondary to breast cancer: how choice of measure influences diagnosis, prevalence, and identifiable risk factors. *Lymphology.* Mar 2008; 41(1): 18-28. PMID 18581955
7. Whitworth P, Vicini F, Valente SA, et al. Reducing rates of chronic breast cancer-related lymphedema with screening and early intervention: an update of recent data. *J Cancer Surviv.* Aug 10 2022. PMID 35947288
8. Barrio AV, Eaton A, Frazier TG. A Prospective Validation Study of Bioimpedance with Volume Displacement in Early-Stage Breast Cancer Patients at Risk for Lymphedema. *Ann Surg Oncol.* Dec 2015; 22 Suppl 3(0 3): S370-5. PMID 26085222
9. Blaney JM, McCollum G, Lorimer J, et al. Prospective surveillance of breast cancer-related lymphoedema in the first-year post-surgery: feasibility and comparison of screening measures. *Support Care Cancer.* Jun 2015; 23(6): 1549-59. PMID 25398360
10. Ridner SH, Dietrich MS, Boyages J, et al. A Comparison of Bioimpedance Spectroscopy or Tape Measure Triggered Compression Intervention in Chronic Breast Cancer Lymphedema Prevention. *Lymphat Res Biol.* Dec 2022; 20(6): 618-628. PMID 35099283
11. Ridner SH, Dietrich MS, Cowher MS, et al. A Randomized Trial Evaluating Bioimpedance Spectroscopy Versus Tape Measurement for the Prevention of Lymphedema Following Treatment for Breast Cancer: Interim Analysis. *Ann Surg Oncol.* Oct 2019; 26(10): 3250-3259. PMID 31054038
12. Soran A, Ozmen T, McGuire KP, et al. The importance of detection of subclinical lymphedema for the prevention of breast cancer-related clinical lymphedema after axillary lymph node dissection; a prospective observational study. *Lymphat Res Biol.* Dec 2014; 12(4): 289-94. PMID 25495384

13. Laidley A, Anglin B. The Impact of L-Dex(®) Measurements in Assessing Breast Cancer-Related Lymphedema as Part of Routine Clinical Practice. *Front Oncol.* 2016; 6: 192. PMID 27656420
14. Koelmeyer LA, Borotkanics RJ, Alcorso J, et al. Early surveillance is associated with less incidence and severity of breast cancer-related lymphedema compared with a traditional referral model of care. *Cancer.* Mar 15 2019; 125(6): 854-862. PMID 30521080
15. Kilgore LJ, Korentager SS, Hangge AN, et al. Reducing Breast Cancer-Related Lymphedema (BCRL) Through Prospective Surveillance Monitoring Using Bioimpedance Spectroscopy (BIS) and Patient Directed Self-Interventions. *Ann Surg Oncol.* Oct 2018; 25(10): 2948-2952. PMID 29987599
16. Whitworth PW, Cooper A. Reducing chronic breast cancer-related lymphedema utilizing a program of prospective surveillance with bioimpedance spectroscopy. *Breast J.* Jan 2018; 24(1): 62-65. PMID 29063664
17. Erdogan Iyigun Z, Selamoglu D, Alco G, et al. Bioelectrical impedance for detecting and monitoring lymphedema in patients with breast cancer. Preliminary results of the florence nightingale breast study group. *Lymphat Res Biol.* Mar 2015; 13(1): 40-5. PMID 25526543
18. Shah C, Vicini F, Beitsch P, et al. The use of bioimpedance spectroscopy to monitor therapeutic intervention in patients treated for breast cancer related lymphedema. *Lymphology.* Dec 2013; 46(4): 184-92. PMID 25141461
19. Lim SM, Han Y, Kim SI, et al. Utilization of bioelectrical impedance analysis for detection of lymphedema in breast Cancer survivors: a prospective cross sectional study. *BMC Cancer.* Jul 08 2019; 19(1): 669. PMID 31286884
20. Kaufman DI, Shah C, Vicini FA, et al. Utilization of bioimpedance spectroscopy in the prevention of chronic breast cancer-related lymphedema. *Breast Cancer Res Treat.* Dec 2017; 166(3): 809-815. PMID 28831632
21. Whitworth PW, Shah C, Vicini F, et al. Preventing Breast Cancer-Related Lymphedema in High-Risk Patients: The Impact of a Structured Surveillance Protocol Using Bioimpedance Spectroscopy. *Front Oncol.* 2018; 8: 197. PMID 29946531
22. Jeffers EJ, Wagner JL, Korentager SS, et al. Breast Cancer-Related Lymphedema (BCRL) and Bioimpedance Spectroscopy: Long-Term Follow-Up, Surveillance Recommendations, and Multidisciplinary Risk Factors. *Ann Surg Oncol.* Oct 2023; 30(10): 6258-6265. PMID 37535267
23. National Comprehensive Cancer Network. Clinical Practice Guidelines in Oncology: Survivorship. Version 1.2023. Updated March 24, 2023. https://www.nccn.org/professionals/physician_gls/pdf/survivorship.pdf. Accessed November 27, 2023.
24. National Comprehensive Cancer Network. Clinical Practice Guidelines in Oncology: Breast Cancer. Version 4.2023. Updated March 23, 2023. https://www.nccn.org/professionals/physician_gls/pdf/breast.pdf. Accessed November 26, 2023.